



Chaos and Correlation
International Journal, May 8, 2012

Бифуркация энергии связи и хаос в атомных ядрах

Binding energy bifurcation and chaos in atomic nuclei

Alexander P. Truney (Toronto, Canada)

Alexander P. Truney

В работе рассмотрена модель хаотического поведения нуклонов в атомных ядрах, построенная на основе модели ядерных взаимодействий и статистики Ферми-Дирака.

The model of chaotic behavior of nucleons in nuclei, based on the model of nuclear interactions and the Fermi-Dirac statistics is discussed.

Ключевые слова: нейтрон, протон, ядро, ядерная оболочка, хаос, энергия связи

Keywords: Binding Energy, Chaos, Neutron, Nuclei, Nuclei Shell, Proton.

It is known that the binding energy of nucleons in atomic nuclei depends on a regular motion of protons and neutrons in the nuclear shells, and on the chaotic behaviour of nucleons, which correlates with uncertainty in the measurement of the mass of the nuclides [1-3]. The concept of quantum chaos [4-5] is the basic model of chaotic behaviour of the nucleons.

We consider the model of the bifurcation of the binding energy in atomic nuclei, based on the generalized dynamics of the Verhulst-Ricker-Planck equation [6]. To derive the equations of the model the results of the theory of strong interactions of nucleons in nuclei [7-8] used. According to this theory there is a relationship between the size of the nucleus, binding energy and the interaction parameter, which can be written as follows:

$$r_n E = \sqrt{S b_{nl}^A} = \beta(A) A \quad (1)$$

Here, $A = N + Z$ - the number of nucleons (protons + neutrons), as the units used the speed of light, Planck constant and electron mass. The binding energy is

determined by the number of nucleons with a total mass of proton and electron, thus $E = A(m_p / m_e + 1) - M_A / m_e$.

Since equation (1) must be shared with the standard expression of the size of the nucleus, $r(A) = r_0 A^{1/3}$, reflecting the weak compressibility of nuclear matter, we can define the left-hand side of equation (1) using experimental data [9]. As a result, we find the product of the binding energy and nuclei radius depending on the number of nucleons - Fig. 1. For consistency with the data [9], we put

$$\beta(A) = 0.05325 \ln A.$$




Fig. 1: The product of the binding energy and nuclei radius depending on the number of nucleons according to [9].

Using this correlation, we can represent equation (1) as

$$E_A = \beta A / r_A \tag{2}$$

Now we can construct a discrete model of the energy levels in nuclei as follows:

$$E_{A+1} E_A^2 = \frac{\beta(A+1)(A+1)(\beta(A)A)^2}{r_{A+1} r_A^2} = \frac{A}{4\pi r_A^3 / 3} \frac{4\pi r_A}{3r_{A+1}} \beta(A+1)(A+1)(\beta(A)A)^2 \tag{3}$$

On the other hand, the density of nucleons can be related to the binding energy due to Fermi-Dirac statistics, we have

$$n_A = \frac{A}{4\pi r_A^3 / 3} = \frac{g_Z Z / A}{e^{(E_Z - \mu_Z)/\theta} + 1} + \frac{g_N N / A}{e^{(E_N - \mu_N)/\theta} + 1} \quad (4)$$

Here g_i, E_i, μ_i, θ are the weight factors, energy and chemical potential of protons and neutrons, and the statistical temperature of the nucleon, respectively. Model (3) - (4) was investigated in a wide range of parameters. Let us consider the results obtained in the simplified model under the condition of equality of chemical potentials and energy of protons and neutrons

$$\mu_N = \mu_Z = \mu_A = \theta \ln \alpha, \quad E_N = E_Z = -E_A / A.$$

In this case, the model can be written as

$$\begin{aligned} x_{A+1} x_A^2 &= \frac{K(1 + 1/A)^{2/3} \beta^2(A) \beta(A+1)}{e^{-x_A} + \alpha} \\ x_A &= -\frac{E_A}{A\theta}, \quad K = \frac{4\pi\alpha g_A}{3A\theta^3} \\ g_A &= g_N + g_Z; \quad \beta(A) = 0.05325 \ln A \end{aligned} \quad (5)$$

To close the model (5), it is necessary to formulate the law of temperature and the weight factor change with the number of nucleons. We use a simple hypothesis, which follows from the theory of the Fermi gas of elementary particles [10] that these parameters are proportional to the cube of the boundary momentum, which in turn is determined by the size of the system:

$$\theta = k_1 p_0^3, \quad g_A = k_2 p_0^3, \quad p_0^2 = k_3 / r_A \quad (6)$$

Hence, we find that the temperature and the weight factor decreases with increasing number of nucleons as follows

$$\theta = T_0 A^{-1/2}, \quad g_A = g_0 A^{-1/2} \quad (7)$$

Under conditions (6)-(7), the parameter K on the right side of equation (5) does not depend on the number of nucleons. Let us consider the behaviour of the chemical potential depending on the number of nucleons. Above we assume that the chemical potentials of protons and nucleons are equal and, moreover, their

relation to temperature is a constant, which coincides with the logarithm of the fine structure constant. To test this hypothesis, let consider functions

$$f(A) = -\mu_A / \theta \ln 137$$

$$f_0(T_0) = \sum_{A \geq 12} f(A) / \sum_{A \geq 12} A(N, Z) \quad (8)$$

Using data [9] and equations (5) - (7), we can calculate functions (8) – see Fig. 2-3. Data [9] plotted in Fig. 2-3 show that the chemical potential of nucleons reaches the theoretical value $\mu_A = \theta \ln \alpha$ for the number of nucleons over 12 and for $T_0 > 20$ MeV . Note that the chemical potential of the bound nucleon system is negative, whereas the chemical potential of free fermions is positive and limited by the Fermi energy at zero temperature - see [10-11].




Fig. 2: Chemical potential over temperature as a function of the number of nucleons, calculated on equations (5) - (7), and data [9]. $f(A) = -\mu_A / \theta \ln 137$.

There is a critical point at $T_0 \approx 20$ MeV as it shown in Figure 3. We suggest that real nuclides have a temperature over critical temperature. Therefore a constant in eq. (7) determined and a linear dependence of chemical potential and temperature established.




Figure 3: The chemical potential parameter as a function of T_0 .




Figure 4: Binding energy per nucleon as a function of mass number calculated on eq. (5)-(7) at $\alpha = 1/137$; $K = 0.0371$.

For light nuclei, the chemical potential, as well as other parameters of the model (5)-(7) deviates from the theoretical dependence (6). Nevertheless, we use the model (5), starting with the deuterium nucleus contains two nucleons. We set the starting point at $x_2 = 0.2$. As a result, we find that the structure of energy levels, which is implemented in a system of nucleons with higher temperature $T_0 = 52.858 MeV$ - Figure 4. In this case, the first bifurcation point for the

binding energy of light nuclei corresponds to the carbon isotope ^{12}C , and the second bifurcation point - nickel isotope ^{58}Ni .

With the number of nucleon increasing the energy levels are split series at 2, 4, 8, 16 sublevels, as shown in Figure 4. The specific structure “four rats”, first observed in [6], is formed by increasing parameter K - Figure 5. It was also shown in [6] that there is the transition to chaotic behaviour in a model (5) in the region $\alpha = e^2 / \hbar c \leq 1/137$. It looks like the fine structure constant could be calculated from model (5) as a transition point between regular and chaotic behaviour of nucleons in a nuclei.

It was established that the transition to chaotic behaviour in a model (5) is also observed in violation of the equality of chemical potentials of the two kinds of nucleons – Figure 6. If the chemical potentials of protons and neutrons are strong differ, than the structure shown in Figure 7 forming, which superficially similar to the experimental dependence - Fig. 8.




Figure 5: Binding energy per nucleon as a function of mass number calculated on eq. (5)-(7) at $\alpha = 1/137$; $K = 0.0839$.

Let us give an interpretation of the results. Model (3) - (7) is a thermodynamic one. It shows how the binding energy changing if one nucleon in the nuclei added, taking into account changes in density according to the Fermi-

Dirac distribution at finite temperature. It is well known that the binding energy of nucleons in the nucleus depends on the number of neutrons and protons. Standard semi-empirical formula describing the binding energy is given by [11]

$$E_b = a_1 A - a_2 A^{2/3} - a_3 Z(Z-1)A^{-1/3} - a_4 (N-Z)^2 A^{-1} + a_5 A^{-3/4} \quad (9)$$

$$a_1 = 14; a_2 = 13; a_3 = 0.585; a_4 = 19.3; a_5 = 33\delta(A, N, Z)$$

Here are shown current values of the coefficients derived from data [9]. All coefficients are given in MeV. In this expression, the function $\delta(A, N, Z)$ is defined as:

$$\delta = 1 \text{ for even } Z, N;$$

$$\delta = -1 \text{ for odd } Z, N;$$

$$\delta = 0 \text{ in all other cases.}$$

The first and fourth term on the right side of expression (9) depend on the kinetic energy of nucleons, which is calculated on the basis of statistics (4) at zero temperature [11]. However, the data in Fig. 2 and eq. (6) - (8) show that temperature not zero and the chemical potential can be varied with temperature by other way than theory of the Fermi gas of free particles predicts, like it explained in [10-11] and other university books. In particular, the chemical potential in a system of nucleons in nuclei is negative, as well as the binding energy.

There is a minimal constant $T_0 \approx 21 \text{ MeV}$ for which is still running a linear relationship of temperature and chemical potential in the area $A > 12$ - Fig. 2. Consider the solution of equation (5) in the case $T_0 = 21 \text{ MeV}$ - Fig. 8. There are two bifurcation points $A = 25; 148$, between which the calculated curve fits data [9]. One branch of the solution diverges in the region $A > 300$ corresponding not yet open nuclei with many nucleons, while the other branch comes to zero. We can assume that the real nuclei temperature constant $T_0 \approx 21 \text{ MeV}$, which agrees with the value of the constants $a_4 = 19.3 \text{ MeV}$ in the semi-empirical equation (9). Further studies will show whether it is possible to predict binding energy based on model (3)-(4) with accuracy exceeding the semi-empirical equation (9).




Figure 6: Binding energy per nucleon as a function of mass number calculated on eq. (5)-(7) at $K = 0.063; \mu_p / \theta = -\ln(137); \mu_n / \theta = -\ln(171)$.




Figure 7: Binding energy per nucleon as a function of mass number calculated on eq. (5)-(7) at $K = 0.07; \mu_p / \theta = -\ln 137; \mu_n / \theta = -2\ln 137$.




Figure 8: Binding energy per nucleon as a function of mass number calculated on eq. (5)-(7) at $T_0 = 21\text{MeV}$; $K = 0.03623$.

Equation (5) is the simplest model describing the dynamics associated with changes in the number of fermions particles. The obtained results on the chaotic behaviour of the nucleons in the bound system indicate the complexity of describing the state of the nuclei, since the splitting of energy levels can occur not only due to the dynamic conditions imposed by the presence of nuclear interaction and the orbital motion, but also because of statistical reasons related to the influence of temperature in accordance with statistics of fermions.

References

1. P. Leboeuf. Regularity and chaos in the nuclear masses/ Lect. Notes Phys. **652**, Springer, Berlin Heidelberg 2005, p.245, J. M. Arias and M. Lozano (Eds.).
2. Jorge G. Hirsch, Alejandro Frank, Jose Barea, Piet Van Isacker, Victor Velazquez. Bounds on the presence of quantum chaos in nuclear masses//Eur. Phys. J. A 25S1 (2005) 75-78
3. Jose Barea, Alejandro Frank, Jorge G. Hirsch, Piet Van Isacker. Nuclear masses set bounds on quantum chaos// Phys.Rev.Lett. 94 (2005) 102501
4. Luca Salasnich. Chaos and Quantum Chaos in Nuclear Systems/ In 6-th workshop "Perspectives on Theoretical Nuclear Physics", Cortona (Italy), 12-14 October 1995
5. E. Caurier, J.M.G. Gomez, V.R. Manfredi, L. Salasnich. Quantum Chaos in $A=46-50$ Atomic Nuclei// Phys. Lett. B365 (1996) 7.

6. Volov D.B. The generalized Verhulst-Ricker-Planck dynamics and its relation to the fine-structure constant. Bulletin of Volga Region Transportation. # 5 (29). 82-90. 2011. Д.Б. <http://www.sciteclibrary.ru/rus/catalog/pages/11612.html>
7. А. Р. Trunев. The structure of atomic nuclei in Kaluza-Klein theory // Политематический сетевой электронный научный журнал Кубанского государственного аграрного университета (Научный журнал КубГАУ) [Электронный ресурс]. – Краснодар: КубГАУ, 2012. – №02(76). С. 862 – 881. – Режим доступа: <http://ej.kubagro.ru/2012/02/pdf/70.pdf>
8. Alexander P. Trunев. Nuclei shells and periodic trends//Chaos and Correlation, April 19, 2012, http://chaosandcorrelation.org/Chaos/CR_1_4_2012.pdf
9. JAGDISH K. TULI. NUCLEAR WALLET CARDS (Seventh edition). April 2005, NATIONAL NUCLEAR DATA CENTER, www.nndc.bnl.gov
10. Ландау Л.Д., Лифшиц Е.М. Теоретическая физика: Т.5. Статистическая физика. Ч.1. – М., Наука. 1976. – 584 с.
11. Marcelo Alonso, Edward J. Finn. Fundamental University Physics. III Quantum and Statistical Physics. – Addison-Wesley Publishing Company, 1975.